REFERENCES:

1. Sharma D, Mittal R, Gupta A, Singh K, Nair A, (2010), Quantitative Bioanalysis by LC-MS/MS: A Review; journal of pharmaceutical and biomedical sciences, 7 (01),1-9.
2. Rajpoot B,(2011), Validation and process development: A review. International. Research. Journal of Pharmacy, 2(1), 32-39.
3. Braggio S, Barnaby R, Grossi P, Cugola M, (1996), a strategy for validation methods, Journal of Pharmaceutical and Biomedical Analysis, 14,375-388.
4. Sethi P,(2001), High performance chromatography: quantitative analytical pharmaceutical formulations. 1st edition, Chapter 1. Introduction. Published by CBS publishers new Delhi, 3-40.
5. Wal P, Kumar B, Bhandari A, Rai A, Wal A,(2010), Bioanalytical method development-Determination of drugs in biological fluids. Journal of Pharmaceutical. Sciences and. Technology, 2(10), 333-347.
6. Xiaofang F, Yiping L and Huwei L, (2005), Sample preparation for pharmaceutical analysis. Analytical and Bioanalytical Chemistry, 381, Page no. 75–77.
7. Poole C, (2003), New trends in solid-phase extraction. Trend. Analytical Chemistry, 22(6), 362-373.
8. Yang L, Jensen D, Vail M, Dadson A, Linford M, (2010), Direct modification of hydrogen/deuterium-terminated diamond particles with polymers to form reserved and strong cation exchange solid phase extraction sorbents. Journal of Chromatography A, 1217, 7621-7629.
9. Kole P, (2011), Recent advances in sample preparation techniques for effective bioanalytical methods, Biomedical Chromatography, 199–217,
10. US Food and Drug Administration, Guidance for industry- Bioanalytical method validation, Center for Drug Evaluation and Research, Rockville, MD, 2001.
11. Health Canada, Guidance for industry- Conduct and analysis of Bioavailability and Bioequivalence studies – Part A: Oral dosage formulations used for systemic effects, authority of the minister of health, 1992.
12. Dofestefano A, Bansal S, (2007), Key elements of bioanalytical method validation for small molecules. Journal of the American Association of Pharmaceutical Scientists, 9(1), E109-E114.

13. Kalakuntla R, Kumar K, (2009), Bioanalytical method validation: A quality assurance auditor view point. Journal Pharmaceutical. Sciences and Research, 1(3), 1-10.
14. Nicola C, Ernest Y, Wong K, Juan F and Bajaj N,(2007), Determination of Carryover and Contamination for Mass Spectrometry – Based Chromatographic Assays; Journal of the American Association of Pharmaceutical Scientists, 9(3) Article 42.
15. Côté C, Bergeron A, Mess N, Furtado M, Garofolo F. (2009), Matrix effect elimination during LC-MS/MS bioanalytical method development; Bioanalysis, Oct 1(7), 1243-57.
16. Khan.F, Sagar.D, Vivake.S, Lohiya R, Umekar M, (2010), Development of UV Spectrophotometric method for the simultaneous estimation of Piroxicam and Paracetamol in tablet by simultaneous Equation, Absorbance ratio and Absorbance Correction method;
    Journal of Pharmacy Research, 3(6), 1432-1433.
17. Mallesh K, Abdul B, Rao A, Diwan P, 2011, Estimation of Piroxicam in Proliposomal Formulation Using RP HPLC Method, International Journal of Chemical and Analytical Science, 2(9), 1193-1196.
18. Devkhile A, Shaikh K, (2010), Development and Validation of LC Method for Simultaneous Determination of Piroxicam and Paracetamol in New Pharmaceutical Formulation Der Pharmacia Lettre, 2(5): 237-244.
19. Fengshan Y, Yangbin Z, Fang C and Lanhua C, (2009), Chemiluminescence method for the determination of piroxicam by the enhancement of the tris-(4,7-diphenyl-1,10-phenanthrolinedisulphonic acid) ruthenium(II)(RuBPS)–cerium(IV) system and its application, Luminescence**;** 24, 50–54.
20. Fegade, J, Bhole R, Shaikh I, Chaudhari R, Patil V, (2009), Development and Validation of Reverse Phase High Performance Liquid Chromatographic Method for Simultaneous Estimation of Paracetamol and Piroxicam in Tablet, International Journal of PharmTech Research ,1(2),184-190.
21. Jiang G, Ge D, Wanga J, Xi Z, (1991), A HPLC method for determining piroxicam in body fluidsZhongguo Yao Li Xue Bao., 12(4),381-384.
22. Ji Y, Lee W, Kim H, Jeong W, Lee S, (2005), Simultaneous determination of piroxicam, meloxicam and tenoxicam in human plasma by liquid chromatography with tandem mass spectrometry. Journal of Chromatography B, 826(1-2), 214-9.
23. Li H, Zhang C, Wang J, Jiang Y, Fawcett P, Gu J, (2010), Simultaneous quantitation of paracetamol, caffeine, pseudoephedrine, chlorpheniramine and cloperastine in human

plasma by liquid chromatography-tandem mass spectrometry, Journal of Pharmaceutical and Biomedical Analysis.,51(3),716-22.

24. Hao Z, Shouhong G, Wansheng C, Yanqiang Z, Xuetao J and Yuanying P, (2008), Simultaneous Quantitation of Paracetamol, Pseudoephedrine and Chlorpheniramine in Dog Plasma by LC-MS-MS, Chromatographia , 68(3-4), 251-257.
25. Qing L, Lin S, Yuan G, Lin Z, Yang G, Chun C, Qiao C, Xiao Q, (2011) , Quantification of acetaminophen (paracetamol) in human plasma and urine by stable isotope-dilution GC-MS and GC-MS/MS as pentafluorobenzyl ether derivative, Journal Of Chromatography B Analytical Technologies In The Biomedical And Life Sciences ,879(7-8), 461-466.
26. Levent A, (2002), HPLC Method for the Analysis of Paracetamol, Caffeine and Dipyrone, Turk Journal of Chemistry, 26, 521 - 528.
27. Reddy B and Reddy M, (2009), RP-HPLC Method for Simultaneous Estimation of Paracetamol and Ibuprofen in Tablets, Asian Journal of Research Chemistry, 2(1), 70-72.
28. Khan H, Ali M, Ahmad S, Niyaz A , Ahuja A, Baboota S, (2012), validated uplc/q-tof-ms method for simultaneous determination of aceclofenac, paracetamol, and their degradation products in tablets, Journal of Liquid Chromatography & Related Technologies ,35(1),109-128.
29. Hong-gang L, Hong Y, Zou-rong R, Bo J, (2010), Simultaneousdetermination of paracetamol, pseudoephedrine, dextrophan and chlorpheniramine in human plasma by liquid chromatography–tandem mass spectrometry, Journal of Chromatography B, 878 (7–80), Pages 682–688.
30. Pattan S, Jamdar S, Godge R, Dighe N, Daithankar A, (2009), RP- HPLC Method for Simultaneous Estimation of Paracetamol and Etoricoxib from Bulk and Tablets, Journal of Chemical and Pharmaceutical Research, 1(1): 329-335.
31. Ophelia P, Sherry S, Moses S, Chow S, (2005), Simultaneous determination of paracetamol and dextropropoxyphene in human plasma by liquid Chromatography/tandem mass spectrometry: application to clinical bioequivalence studies, Rapid Communications in Mass Spectrometry,19(6), 767–774.

32. Dattatray T, Handa T, Shah R and Saranjit S, (2011), Successful characterization of degradation products of drugs using LC-MS tools: Application to piroxicam and meloxicam, , Journal of Analytical Methods,3, 2864-2872.
33. Małgorzata S, Jan K, Tarsa M and Marek Ż,(2009), Determination of Piroxicam and Degradation Products in Drugs by TLC, Chromatographia , 69,(3-4), 351-356.
34. Marın A, Barbas C, (2004), a LC/MS for the degradation profiling of cough–cold products under forced conditions, Journal of Pharmaceutical and Biomedical Analysis, 1035–1045.
35. Micaela C, Jan N, Fidder A, Albert G, Daan T, (2007), Liquid Chromatography/Tandem Mass Spectrometry Detection of Covalent Binding of Acetaminophen to Human Serum Albumin , The American Society for Pharmacology and Experimental Therapeutics **,**35 (8),1408-1417.
36. Qin-you T, Rong-hua Z, Huan-de L, Feng W, Miao Y, Li-bo D, (2012), Simultaneous quantitative determination of paracetamol and its glucuronide conjugate in human plasma and urine by liquid chromatography coupled to electrospray tandem mass spectrometry, Application to a clinical pharmacokinetic study, Journal of Chromatography B, 893–894, 162–167.
37. Gondalia R, Mashru R, Savaliya P, **(**2010), Development and Validation of Spectrophotometric Methods for Simultaneous Estimation of ibuprofen and paracetamol in Soft gelatin capsule by Simultaneous Equation Method International Journal of ChemTech Research,.2(4),1881-1885.
38. Celma C, Allue J, Prunonosa J., Peraire C, Obach R, (2000), Simultaneous determination of paracetamol and chlorpheniramine in human plasma by liquid chromatography-tandem mass spectrometry, Journal of Chromatography A, 870(1), 77-86.
39. Hewavitharana A , Lee S, Dawson P , Markovich D, Shaw P, (2008), Development of an HPLC–MS/MS method for the selective determination of paracetamol metabolites in mouse urine, Analytical,374 (1), 106–111.
40. Khade T, Saindane D and Pore Y, (2010), Simultaneous Spectrophotometric Determination of Etoricoxib and Paracetamol in a Laboratory Mixture, International Journal of PharmTech Research, 2(1), 623-626.